

STUDIES IN NATURE CONSERVATION

J.B. Kirkpatrick BA (Hons), PhD *University of Melbourne* AM

Submitted in fulfilment of the requirements for the degree of

Doctor of Science

University of Tasmania

Hobart

July 2005

VOLUME I

Cent
Thesis
KIRK PATRICK
D. Sc.
2005
Vol. 1

STATEMENT OF ORIGINALITY

This volume contains a substantial part of my research contribution to the conservation of nature. It consists 110 refereed papers or book chapters, one invited editorial and one chapter from a sole-authored book. None of this work has been submitted by me as part of a thesis for any other degree, although many papers have been adapted from theses written by the many honours and postgraduate students I have supervised since 1973 (Table 1).

I am sole author of 22 of the writings included in this volume. As can be gauged from Table 1, which outlines the contributions of the various authors of papers contained herein, and from the acknowledgements in the papers, I have benefited from working with many outstanding research students and colleagues, among them: Kerry Bridle, David Bowman, Mick Brown, Kath Dickinson, Rod Fensham, Neil Gibson, Louise Gilfedder, Stephen Harris, Chris Harwood, Tony Moscal and Jenny Scott. I could not have undertaken this work without the help of the general staff of the School of Geography and Environmental Studies and its predecessor academic units.

I declare the above information to be true

kpatrick 25 July 2005




Table 1. Contribution to papers with multiple authors. Estimated percentages are shown. n.a. = not applicable, * paper developed from the thesis of a student co-author who I supervised. + see table of contents for paper codes.

Paper+	Conception	Data collection	Data analysis	Writing up
2.2	50	50	90	95
2.3	60	50	90	95
2.5	50	20	30	30
3.1.3	100	50	100	95
3.1.4	50	50	80	95
3.1.6	100	50	70	95
3.1.7*	40	0	10	50
3.1.8	100	50	80	90
3.1.9	100	0	50	80
3.1.10*	50	5	10	50
3.1.11*	40	5	10	50
3.1.12*	40	5	10	50
3.1.13*	40	0	0	50
3.1.14*	60	0	20	60
3.1.15*	50	5	10	60
3.1.16	100	50	100	95
3.1.17	60	0	100	80
3.1.18	25	0	100	80
3.2.2*	60	5	70	80
3.2.3	70	50	70	90
3.2.4*	50	0	0	50
3.2.5*	30	0	0	50
3.2.6*	30	0	0	50
3.2.7*	30	0	0	50
3.2.8*	50	20	20	50
3.2.9	70	5	50	60
3.2.10	100	30	50	95
3.2.11*	70	0	10	20
3.2.12*	70	5	10	50
3.3.1	50	50	100	95
3.3.2	50	70	100	95
3.3.3	50	50	100	95
3.3.4	50	30	70	95
3.4.1	70	50	100	95
3.4.2	70	50	100	95
3.5.1	50	5	100	95
3.5.2*	50	5	5	50
3.5.3*	50	0	20	50
3.5.4*	50	0	10	50

Paper	Conception	Data collection	Data analysis	Writing up
3.5.5	100	20	80	95
3.5.6*	50	10	30	50
4.2	50	0	100	95
4.3*	50	0	20	50
4.4*	40	0	10	50
4.5*	40	0	10	50
4.6*	40	0	10	50
4.7*	40	0	10	50
4.8	50	0	80	95
4.9	50	0	80	95
4.10	40	0	50	60
4.11*	40	5	20	50
4.12*	40	5	10	50
4.13*	50	0	50	50
5.5*	70	20	80	60
5.6*	30	20	30	10
5.7*	60	0	50	70
6.1.1	100	50	100	95
6.1.3	50	0	80	95
6.1.5	50	0	100	95
6.1.6*	60	0	40	50
6.1.7*	40	0	40	40
6.1.8	60	0	50	30
6.2.1	50	na	na	90
6.2.2*	40	0	40	50
6.2.3*	50	0	30	50
6.2.4*	50	0	30	50
6.3.1	80	0	100	95
6.3.2	80	30	100	95
6.3.3*	25	0	0	20
6.3.4*	50	na	na	50
6.4.1	50	50	40	50
6.4.2*	40	0	30	50
6.4.3*	40	30	30	50
6.4.4*	20	0	30	30
6.4.5	60	0	100	95
6.4.6*	30	0	30	20
6.4.7	80	20	100	95
6.5.2*	50	5	50	50
6.5.3*	50	5	30	40
6.5.4	50	50	50	50
6.5.5	80	0	90	80
6.5.6	70	0	80	80
7.1*	70	na	20	30

Paper	Conception	Data collection	Data analysis	Writing up
7.2	100	60	100	95
7.5	80	40	80	70
7.7	100	50	90	95
7.9	50	50	50	80
7.10	90	50	80	95
7.12*	70	0	10	20

TABLE OF CONTENTS

Section	Page
Introduction	1
Papers	
1. Research approach	
1.1 Kirkpatrick, J.B., 2000. The political ecology of biogeography. <i>Journal of Biogeography</i> 27, 45-48.	12
1.2. Kirkpatrick, J.B., 1997. The future of Australian plant ecology - incisively idiographic or nebulously nomothetic? In Klomp, N. and Lunt, I. (eds.). <i>Frontiers in Ecology</i> . Elsevier Science, Oxford, 289-298.	16
1.3. Kirkpatrick, J.B., 1999. The politics of the media and ecological ethics. In Wills, R. and Hobbs, R. (eds.) <i>Ecology for Everyone</i> . Surrey-Beatty, Sydney, 36-41.	26
2. Floristic biogeography of Tasmania	
2.1. Kirkpatrick, J.B., 1982. Phytogeographical analysis of Tasmanian alpine floras. <i>Journal of Biogeography</i> 9, 255-271.	33
2.2. Kirkpatrick, J.B. and Brown, M.J., 1984. Numerical analysis of Tasmanian higher plant endemism. <i>Botanical Journal of the Linnaean Society of London</i> 88, 165-183.	50
2.3. Kirkpatrick, J.B. and Brown, M.J., 1984. The palaeogeographic significance of local endemism in Tasmanian higher plants. <i>Search</i> 15, 112-113.	69

2.4. Kirkpatrick, J.B., 1986. Tasmanian alpine biogeography and ecology and interpretation of the past. In Barlow, B.A. (ed.). *Flora and Fauna of Alpine Australasia: Ages and Origins*. CSIRO, Melbourne, pp. 229-242. 71

2.5. Nunez, M., Kirkpatrick, J.B. and Nilsson, C., 1996. Rainfall estimation in south west Tasmania using satellite images and phytosociological calibration. *International Journal of Remote Sensing* 17, 1583-1600. 85

2.6. Kirkpatrick, J.B., 2002. Factors influencing the spatial restriction of vascular plant species in the alpine archipelagoes of Australia. In C. Körner & E.M. Spehn (eds.): *Mountain Biodiversity: a Global Assessment*. Parthenon Publishing, London, pp. 155-164. 103

3. Ecology of natural vegetation

3.1. Cold climate vegetation

3.1.1. Kirkpatrick, J.B., 1983. Treeless plant communities of the Tasmanian high country. *Proceedings of the Ecological Society of Australia* 12, 61-77. 113

3.1.2. Kirkpatrick, J.B. 1989. The comparative ecology of mainland Australian and Tasmanian alpine vegetation . In R. Good (ed.) *The Scientific Significance of the Australian Alps*. Australian Academy of Science, Canberra, 127-142. 130

3.1.3. Kirkpatrick, J.B. and Bridle, K.L., 1998. Environmental relationships of floristic variation in the alpine vegetation of south-east Australia. *Journal of Vegetation Science* 9, 251-260. 146

- 3.1.4. Kirkpatrick, J.B. and Harwood, C.E., 1980. The vegetation of an infrequently burned Tasmanian mountain region. *Proceedings of the Royal Society of Victoria* 91, 79-107. 156
- 3.1.5. Kirkpatrick, J.B., 1984. Altitudinal and successional variation in the vegetation of the West Coast Range, Tasmania. *Australian Journal of Ecology* 9, 81-91. 185
- 3.1.6. Kirkpatrick, J.B. and Gibson, N.L., 1984. Dynamics of a Tasmanian bolster heath string fen. *Vegetatio* 58, 71-78. 196
- 3.1.7. Gibson, N. and Kirkpatrick, J.B., 1992. Dynamics of a Tasmanian cushion heath community. *Journal of Vegetation Science* 3, 647-654. 204
- 3.1.8. Kirkpatrick, J.B. and Brown, M.J., 1987. The nature of the transition from sedgeland to alpine vegetation in South West Tasmania: I. Altitudinal vegetation change on four mountains. *Journal of Biogeography* 14, 539-549. 212
- 3.1.9. Kirkpatrick, J.B., Nunez, M., Bridle, K. and Chladil, M., 1996. Explaining a sharp transition from sedgeland to alpine vegetation on Mount Sprent, south-west Tasmania. *Journal of Vegetation Science* 7, 763-768. 223
- 3.1.10. Bridle, K. and Kirkpatrick, J.B., 1997. Local environmental correlates of variability in the organic soils of moorland and alpine vegetation, Mt Sprent, Tasmania. *Australian Journal of Ecology* 22, 196-205. 229
- 3.1.11. Cullen, P.C. and Kirkpatrick, J.B., 1988. The ecology of *Athrotaxis* D. Don (Taxodiaceae) I. Stand structure and regeneration of *A. cupressoides*. *Australian Journal of Botany* 36, 547-560. 239

- 3.1.12. Cullen, P.C. and Kirkpatrick, J.B., 1988. The ecology of *Athrotaxis* D. Don (Taxodiaceae) II. The distributions and ecological differentiation of *A. cupressoides* and *A. selaginoides*. *Australian Journal of Botany* 36, 561-573. 253
- 3.1.13. Whinam, J. and Kirkpatrick, J.B., 1995. Successional sequences in two Tasmanian valley *Sphagnum* peatlands. *Journal of Vegetation Science* 6, 675-682. 266
- 3.1.14. Pharo, E. and Kirkpatrick, J.B., 1994. Vegetation of the alpine sand dunes at Lake Augusta, Tasmania. *Australian Journal of Ecology* 19, 319-327. 274
- 3.1.15. Lynch, A.J.J. and Kirkpatrick, J.B., 1995. Pattern and process in alpine vegetation and landforms at Hill One, Southern Range, Tasmania. *Australian Journal of Botany* 43, 537-554. 283
- 3.1.16. Kirkpatrick, J.B., Bridle, K. and Lynch, A.J.J., 2002. Changes in alpine vegetation related to geomorphological processes and climate change on Hill One, Southern Range, Tasmania, 1989-2000. *Australian Journal of Botany* 50, 753-759. 299
- 3.1.17. Kirkpatrick, J.B. and Scott, J.J., 2002. Change in undisturbed vegetation on the coastal slopes of Subantarctic Macquarie Island 1980-1995. *Arctic, Antarctic and Alpine Research*, 34, 300-307. 306
- 3.1.18. Scott, J.J. and Kirkpatrick, J.B., in press. Changes in Subantarctic Heard Island vegetation at sites occupied by *Poa annua*, 1987-2000. *Arctic, Antarctic and Alpine Research*. 314

3.2 *Eucalypt-dominated vegetation*

- 3.2.1. Kirkpatrick, J.B., 1997. Vascular plant-eucalypt interactions. In. Williams, J. E. and Woinarski, J.C.Z. (eds.). *Eucalypt Ecology: Individuals to Ecosystems*. Cambridge University Press, Cambridge, pp. 227-245. 329
- 3.2.2. Hogg, A. and Kirkpatrick, J.B., 1974. The phytosociology and synecology of some southern Tasmanian eucalypt forests and woodlands. *Journal of Biogeography* 1, 227-245. 348
- 3.2.3. Kirkpatrick, J.B. and Nunez, M., 1980. Vegetation-radiation relationships in mountainous terrain: eucalypt-dominated vegetation in the Risdon Hills, Tasmania. *Journal of Biogeography* 7, 197-208. 367
- 3.2.4. Bowman, D.M.J.S. and Kirkpatrick, J.B., 1984. Geographic variation in the demographic structure of stands of *Eucalyptus delegatensis* R.T. Baker on dolerite in Tasmania. *Journal of Biogeography* 11, 427-437. 379
- 3.2.5. Bowman, D.M.J.S. and Kirkpatrick, J.B., 1986. The establishment, suppression and growth of *Eucalyptus delegatensis* R.T. Baker in multi-aged forests. I. The effects of fire on mortality and seedling establishment. *Australian Journal of Botany* 34, 63-72. 390
- 3.2.6. Bowman, D.M.J.S. and Kirkpatrick, J.B., 1986. The establishment, suppression and growth of *Eucalyptus delegatensis* R.T. Baker in multi-aged forests. II. Sapling growth and its environmental correlates. *Australian Journal of Botany* 34, 73-80. 400
- 3.2.7. Bowman, D.M.J.S. and Kirkpatrick, J.B., 1986. The establishment, suppression and growth of *Eucalyptus delegatensis* in multi-aged forests. III. Intraspecific

allelopathy, competition between adult and juvenile for moisture and nutrients, and frost damage to seedlings. *Australian Journal of Botany* 34, 81-94. 408

3.2.8. Dickinson, K.J.M. and Kirkpatrick, J.B., 1985. The flammability and energy content of some important plant species and fuel components in the forests of southeastern Tasmania. *Journal of Biogeography* 12, 121-134. 422

3.2.9. Fensham, R.J. and Kirkpatrick, J.B., 1992. The eucalypt forest-grassland/grassy woodland boundary in central Tasmania. *Australian Journal of Botany* 40, 123-138. 436

3.2.10. Kirkpatrick, J.B. and Gibson, N., 1999. Towards an explanation of the altitudinal distributions of three species of *Eucalyptus* in central Tasmania. *Australian Journal of Ecology* 24, 123-131. 452

3.2.11. Pyke, A. and Kirkpatrick, J.B., 1994. Growth rate and basal area response curves of four *Eucalyptus* species on Mt Wellington, Tasmania. *Journal of Vegetation Science* 5, 13-24. 461

3.2.12. McKenny, H. and Kirkpatrick, J.B., 1999. The role of fallen logs in the regeneration of tree species in Tasmanian mixed forest. *Australian Journal of Botany* 47, 745-753. 473

3.3 Tropical vegetation

3.3.1. Kirkpatrick, J.B. and Hassall, D.C., 1981. Vegetation of the Sigatoka sand dunes, Fiji. *New Zealand Journal of Botany* 19, 285-297. 480

3.3.2. Kirkpatrick, J.B. and Hassall, D.C., 1985. The vegetation and flora along an altitudinal transect through tropical forest at Mount Korobaba, Fiji. *New Zealand Journal of Botany* 23, 33-46. 493

- 3.3.3. Hassall, D.C. and Kirkpatrick, J.B., 1985. The diagnostic value and host relationships of the dependent synusia in the forests of Mount Korobaba, Fiji. *New Zealand Journal of Botany* 23, 47-54. 507
- 3.3.4. Kirkpatrick, J.B., Fensham, R., Nunez, M. and Bowman, D.M.J.S., 1988. Vegetation-radiation relationships in the wet-dry tropics: granite hills in northern Australia. *Vegetatio* 76, 103-112. 515
- 3.4. Californian coastal sage scrub**
- 3.4.1. Kirkpatrick, J.B. and Hutchinson, C.F., 1977. The community composition of Californian coastal sage scrub. *Vegetatio* 35, 21-33. 525
- 3.4.2. Kirkpatrick, J.B. and Hutchinson, C.F., 1980. The environmental relationships of Californian coastal sage scrub and some of its component communities and species. *Journal of Biogeography* 7, 23-38. 538
- 3.5. Miscellaneous Tasmanian vegetation types**
- 3.5.1. Kirkpatrick, J.B. and Harwood, C.E., 1983. Plant communities of Tasmanian wetlands. *Australian Journal of Botany* 31, 437-451. 554
- 3.5.2. Daley, E.A. and Kirkpatrick, J.B., 2004. Native riparian vegetation in Tasmania. *Cunninghamiana* 8, 409-430. 567
- 3.5.3. Coates, F. and Kirkpatrick, J.B., 1992. Environmental relationships and ecological responses of some higher plant species on rock cliffs in northern Tasmania. *Australian Journal of Ecology* 17, 441-449. 589

- 3.5.4. Walsh, D.A., Kirkpatrick, J.B. and Skira, I.J., 1997. Vegetation patterns, environmental correlates and vegetation change in a *Puffinus tenuirostris* breeding colony at Cape Queen Elizabeth, Tasmania. *Australian Journal of Botany* 45, 71-79. 598
- 3.5.5. Kirkpatrick, J.B., Leaman, T., Pennington, A. and Pollard, T., 2004. Characteristics and dynamics of the sand dune vegetation at North Bay, Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* 138, 37-45. 607
- 3.5.6. Harris, S. and Kirkpatrick, J.B., 1991. The distributions, dynamics and ecological differentiation of *Callitris* species in Tasmania. *Australian Journal of Botany* 39, 187-202. 616
- 4. The conservation ecology of rare or threatened species**
- 4.1. Kirkpatrick, J.B. 1999. *A Continent Transformed- Human Impact on the Natural Vegetation of Australia*. 2nd edition, Oxford University Press, Sydney, chapter 9. 632
- 4.2. Kirkpatrick, J.B. and Gilfedder, L., 1995. Maintaining integrity compared with maintaining rare and threatened taxa in remnant bushland in subhumid Tasmania. *Biological Conservation* 74, 1-8. 644
- 4.3. Woolley, A. and Kirkpatrick, J.B., 1999. Factors related to condition and rare and threatened species occurrence in lowland, humid basalt remnants in northern Tasmania. *Biological Conservation* 87, 131-42. 652
- 4.4. Gilfedder, L. and Kirkpatrick, J.B., 1993. Germinable soil seed and competitive relationships between a rare native species and exotics in a semi-natural pasture in the Midlands, Tasmania. *Biological Conservation* 64, 113-119. 664

- 4.5. Gilfedder, L. and Kirkpatrick, J.B., 1994. Culturally induced rarity? - the past and present distributions of *Leucochrysum albicans* in Tasmania. *Australian Journal of Botany* 42, 404-416. 671
- 4.6. Gilfedder, L. and Kirkpatrick, J.B., 1994. Climate, grazing and disturbance and the population dynamics of *Leucochrysum albicans* at Ross, Tasmania. *Australian Journal of Botany* 42, 417-430. 683
- 4.7. Gilfedder, L. and Kirkpatrick, J.B., 1994. Genecological variation in the germination, growth and morphology of four populations of a Tasmanian endangered perennial daisy, *Leucochrysum albicans*. *Australian Journal of Botany* 42, 431-440. 697
- 4.8. Gilfedder, L. and Kirkpatrick, J.B., 1998. Distribution, disturbance tolerance and conservation of *Stackhousia gunnii* Hook. f. in Tasmania. *Australian Journal of Botany* 46, 1-13. 707
- 4.9. Kirkpatrick, J.B. and Gilfedder, L., 1998. Conserving weedy natives: two Tasmanian endangered herbs in the Brassicaceae. *Australian Journal of Ecology* 23, 466-473. 718
- 4.10. Lynch, A.J.J., Gilfedder, L.G. and Kirkpatrick, J.B., 1999. The Tasmanian endemic shrub, *Acacia axillaris*: conservation ecology applied to the question of rarity or vulnerability. *Australian Journal of Botany* 47, 97-109. 726
- 4.11. Coates, F., Kirkpatrick, J.B. and Minchin, P.R., 1999. Towards an explanation of the causes of the rarity of two Tasmanian *Spyridium* species. *Australian Journal of Ecology* 24, 11-17. 739
- 4.12. Coates, F. and Kirkpatrick, J.B., 1999. Is geographic range correlated with climatic range in Australian *Spyridium* taxa? *Australian Journal of Botany* 47, 755-767. 746

- 4.13. Leeson, K. and Kirkpatrick, J.B., 2004. Ecological and physiological explanations for the restriction of a Tasmanian species of *Ozothamnus* to a single population. *Australian Journal of Botany* 52, 39-45. 759

5. Urban ecology

- 5.1. Kirkpatrick, J.B., 1974. Plant invasion and extinction in a suburban coastal reserve. *Australian Geographical Studies* 12, 107-118. 766
- 5.2. Kirkpatrick, J.B., 1975. Vegetation change in a suburban coastal reserve. *Australian Geographical Studies* 13, 137-153. 778
- 5.3. Kirkpatrick, J.B., 1986. The viability of bush in cities - ten years of change in an urban grassy woodland. *Australian Journal of Botany* 34, 691-708. 795
- 5.4. Kirkpatrick, J.B., 2004. Vegetation change in an urban grassy woodland 1974-2000. *Australian Journal of Botany* 52, 597-608. 813
- 5.5. O'Shea, E.M. and Kirkpatrick, J.B., 2000. The impact of suburbanization on remnant coastal vegetation in Hobart, Tasmania. *Applied Vegetation Science* 3, 243-252. 825
- 5.6. Jellinek, S., Driscoll, D.A., Kirkpatrick, J.B., 2004. Environmental and vegetation variables have a greater influence than habitat fragmentation in structuring lizard communities in remnant urban bushland. *Austral Ecology* 29, 294-304. 835
- 5.7. Zagorski, T., Kirkpatrick, J.B. and Stratford, E. 2004. Gardens and the bush: gardeners attitudes, garden types and invasives. *Australian Geographical Studies* 42, 207-220. 846

6. Human impacts on natural ecosystems

6.1. Clearing

- 6.1.1 Kirkpatrick, J.B. and Dickinson, K.J.M., 1982. Recent destruction of natural vegetation in Tasmania. *Search* 13, 186-187. 860
- 6.1.2. Kirkpatrick, J.B., 1991. The magnitude and significance of land clearance in Tasmania in the 1980s. *Tasforests* 3, 11-14. 862
- 6.1.3. Gilfedder, L. and Kirkpatrick, J.B., 1997. The influence of owners intentions on remnant bush conservation in Tasmania. In P. Hale and D. Lamb (eds.) *Conservation Outside Nature Reserves*. Centre for Conservation Biology, The University of Queensland, Brisbane, pp. 217-225. 866
- 6.1.4. Kirkpatrick, J.B., 1988. Heritage and development in Tasmania. *Australian Geographer* 19, 46-63. 875
- 6.1.5. Gilfedder, L. and Kirkpatrick, J.B., 1998. Factors influencing the integrity of remnant bushland in subhumid Tasmania. *Biological Conservation* 84, 89-96. 893
- 6.1.6. Bowkett, L.A. and Kirkpatrick, J.B., 2003. Ecology and conservation of remnant *Melaleuca ericifolia* stands in the Tamar Valley, Tasmania. *Australian Journal of Botany* 51, 405-413. 901
- 6.1.7. MacDonald, M. and Kirkpatrick, J.B., 2003. Explaining bird species composition and richness in eucalypt-dominated remnants in subhumid Tasmania. *Journal of Biogeography* 30, 1415-1426. 910

- 6.1.8. Pharo, E.J., Kirkpatrick, J.B., Gilfedder, L., Mendel, L. and Turner, P.A.M., in press. Predicting bryophyte diversity in grassland and eucalypt-dominated remnants in subhumid Tasmania. *Journal of Biogeography* 922
- 6.2. Logging**
- 6.2.1. Kirkpatrick, J.B. and Bowman, D.M.J.S., 1982. Clearfelling versus selective logging in uneven-aged eucalypt forests. *Search* 13, 136-141. 959
- 6.2.2. Dickinson, K.J.M. and Kirkpatrick, J.B., 1986. The impact of grazing pressure in clearfelled, burned and undisturbed eucalypt forest. *Vegetatio*, 66, 133-6. 965
- 6.2.3. Dickinson, K.J.M. and Kirkpatrick, J.B., 1987. The short-term effects of clearfelling and slash-burning on the richness, diversity and relative abundance of higher plant species in two types of eucalypt forest on dolerite in Tasmania. *Australian Journal of Botany* 36, 601-16. 969
- 6.2.4. Calais, S.S. and Kirkpatrick, J.B., 1983. Tree species regeneration after logging in temperate rainforest, Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* 117, 77-83. 983
- 6.3. Burning**
- 6.3.1. Kirkpatrick, J.B. and Dickinson, K.J.M., 1984. The impact of fire on Tasmanian alpine vegetation and soils. *Australian Journal of Botany* 32, 613-629. 990
- 6.3.2. Kirkpatrick, J.B., Bridle, K. and Wild, A., 2002. Succession after fire in alpine vegetation on Mount Wellington, Tasmania. *Australian Journal of Botany* 50, 145-154. 1007

- 6.3.3 Thomas, I. and Kirkpatrick, J.B., 1996. The role of coastlines, people and fire in the development of heathlands in northeast Tasmania. *Journal of Biogeography* 23, 717-728. 1017
- 6.3.4. Marsden-Smedley, J. and Kirkpatrick, J.B., 2000. Fire management in Tasmania's Wilderness World Heritage Area: ecosystem restoration using Indigenous-style fire regimes? *Ecological Management and Restoration* 1, 195-203. 1029
- 6.4 Grazing**
- 6.4.1. Gibson, N. and Kirkpatrick, J.B. 1989. Effects of the cessation of grazing on the grasslands and grassy woodlands of the Central Plateau, Tasmania. *Australian Journal of Botany* 37, 55-63. 1038
- 6.4.2. Bridle, K.L. and Kirkpatrick, J.B., 1999. The comparative effects of stock and wild vertebrate grazing on treeless subalpine vegetation, eastern Central Plateau, Tasmania. *Australian Journal of Botany* 47, 817-834. 1047
- 6.4.3. Bridle, K.L., Kirkpatrick, J.B., Cullen, P. and Shepherd, R.R., 2001. Recovery in alpine heath and grassland following burning and grazing, eastern Central Plateau, Tasmania, Australia. *Arctic, Antarctic and Alpine Research* 33, 348-356. 1065
- 6.4.4. Bridle, K.L. and Kirkpatrick, J.B., 2001. Impacts of grazing by vertebrate herbivores on the flower stem production of tall alpine herbs, Eastern Central Plateau, Tasmania. *Australian Journal of Botany* 49, 459-470. 1074
- 6.4.5. Kirkpatrick, J.B., Zacharek, A. and Chappell, K., 2000. Testing methods for mitigation of tree dieback in Tasmanian dry eucalypt forests and woodlands. *Pacific Conservation Biology* 6, 94-101. 1086

6.4.6. Leonard, S. W. J. and Kirkpatrick, J.B., 2004. Effects of grazing management and environmental factors on native grassland and grassy woodland, Northern Midlands, Tasmania. *Australian Journal of Botany* 52, 529-542. 1095

6.4.7. Kirkpatrick, J.B., Gilfedder, L., Bridle, K.L. and Zacharek, A., 2005. The positive and negative conservation impacts of sheep grazing and other disturbances on the vascular plant species and vegetation of lowland subhumid Tasmania. *Ecological Management and Restoration* 6, 51-60. 1109

6.5. Recreational impacts

6.5.1. Kirkpatrick, J.B., 2001. Ecotourism, local and indigenous people, and the conservation of the Tasmanian Wilderness World Heritage Area. *Journal of the Royal Society of New Zealand* 31, 819-829. 1119

6.5.2. Calais, S.S. and Kirkpatrick, J.B., 1986. The impact of trampling on the natural ecosystems of the Cradle Mt. - Lake St. Clair National Park. *Australian Geographer* 17, 6-15. 1130

6.5.3. Whinam, J., Cannell, E.J., Kirkpatrick, J.B. and Comfort, M., 1994. Studies on the potential impact of recreational horseriding on some alpine environments of the Central Plateau, Tasmania. *Journal of Environmental Management* 40, 103-117. 1140

6.5.4. Scott, J.J. and Kirkpatrick, J.B., 1994. Effects of human trampling on the sub-Antarctic vegetation of Macquarie Island. *Polar Record* 30, 207-220. 1155

6.5.5. Bridle, K.L. and Kirkpatrick, J.B., 2003. The impacts of nutrient additions and digging for human waste disposal in natural environments, Tasmania, Australia. *Journal of Environmental Management* 69, 299-306. 1169

6.5.6. Bridle, K.L. and Kirkpatrick, J.B., 2005. An analysis of the breakdown of paper products (toilet paper, tissues and tampons) in natural environments, Tasmania, Australia. *Journal of Environmental Management* 74, 21-30. 1177

7. Nature conservation planning

7.1. Mendel, L.C. and Kirkpatrick, J.B., 2002. Historical progress of biodiversity conservation in the protected-area system of Tasmania, Australia. *Conservation Biology* 16, 1520-1529. 1187

7.2. Kirkpatrick, J.B. and Haney, R.A., 1980. The quantification of developmental wilderness loss - The case of forestry in Tasmania. *Search* 11, 331-335. 1197

7.3. Kirkpatrick, J.B., 1983. An iterative method for establishing priorities for the selection of Nature Reserves: an example from Tasmania. *Biological Conservation* 25, 127-134. 1203

7.4. Kirkpatrick, J.B., 1986. Conservation of plant species, alliances and associations of the treeless high country of Tasmania. *Biological Conservation* 37, 43-57. 1211

7.5. Kirkpatrick, J.B. and Duncan, F., 1987. Tasmanian high altitude grassy vegetation: its distribution, community composition and conservation status. *Australian Journal of Ecology* 12, 73-86. 1227

7.6. Kirkpatrick, J.B., 1990. A synusia-based mapping system for the conservation and management of natural vegetation, with an example from Tasmania, Australia. *Biological Conservation* 53, 93-104. 1242

7.7. Kirkpatrick, J.B., Gilfedder, L., Duncan, F.D. and Harris, S., 1991. Reservation status and priorities for Tasmanian plants 1. Angiospermae (Dicotyledonae). In Banks,

- M.R., Smith, S.J., Orchard, A.E. and Kantvilas, G. (eds). *Aspects of Tasmanian Botany*, Royal Society of Tasmania, Hobart, 163-172. 1253
- 7.8. Kirkpatrick, J.B. and Brown, M.J., 1991. Planning for species conservation. In Margules, C.R. and Austin, M.P. (eds) *Nature Conservation: Cost Effective Biological Surveys and Data Analysis*. CSIRO, Melbourne, 83-89. 1263
- 7.9. Kirkpatrick, J.B. and Brown, M.J., 1994. A comparison of direct and environmental domain approaches to planning reservation of forest higher plant communities and species in Tasmania. *Conservation Biology*, 8, 217-224. 1271
- 7.10. Kirkpatrick, J.B. and Fowler, M., 1998. Locating likely glacial refugia in Tasmania using palynological and ecological information to test alternative climatic models. *Biological Conservation* 85, 171-182. 1279
- 7.11. Kirkpatrick, J.B., 1998. Nature conservation and the regional forest agreement process. *Australian Journal of Environmental Management* 5, 31-37. 1291
- 7.12. Mendel, L.C. and Kirkpatrick, J.B., 1999. Assessing temporal changes in the reservation of the natural aesthetic resource using pictorial content analysis and a grid-based scoring system - the example of Tasmania. *Landscape and Urban Planning* 43, 181-190. 1299

INTRODUCTION

The papers enclosed in this volume were all created with one major motivation, to contribute to the conservation of nature, defining nature in its dualistic sense. In paper 1.1, I described several ways in which biogeographers/ecologists can help create positive outcomes for nature through the research work that they choose to undertake, ranging from revealing interesting stories that might arouse public interest in nature conservation to the provision of a technical and knowledge foundation for conservation measures in matters of public dispute. Paper 1.2 argues the proposition that the particular may be more important than the general in gaining the types of ecological understandings that might enable practical nature conservation. Some of the ethical underpinnings of my research work are discussed in paper 1.3, the main, and unexceptional, point being that ecologists should base their public utterances on an honest reading of their always inadequate data. Attachment to ideas/generalizations/hypotheses can be bad for both science and the living world. The few people who will read this volume will probably pick up some instances of hypothesis drift, where the initial guiding proposition proves less interesting than a byproduct of the research process, and many instances of hypothesis reversal, where an idea that seemed excellent in relation to the data available at the time proves to be incorrect in a context of more data or a wider understanding.

The work in the present thesis was focused largely on Tasmania, a State of islands in the Commonwealth of Australia, constituting approximately 6.9 million ha of land, of which approximately 70% had not been cleared or inundated by 2005. The late Professor Bill Jackson believed that plant ecologists could only be proficient in their trade by knowing an area for decades. Whatever the truth of this proposition, longevity in a place does allow decades scale observations that can often be more powerful than natural or induced experiments in contributing towards an understanding of natural processes (e.g. 6.3.2, 3.1.16; 3.1.17; 5.4). The size and environmental heterogeneity of Tasmania have made it an excellent laboratory for the development and testing of ecological and conservation planning ideas and techniques, many of which have much wider relevance than to one small set of islands.

Before 1970, Barber (1955) and Barber and Jackson (1957) had used observations of genecological variation in *Eucalyptus urnigera* on Mt. Wellington to first postulate the phenomenon of divergence through selection in continuous populations, and Jackson (1968) had created a probabilistic ecosystem model that predicted successional change in lowland western Tasmania in an obscurely published paper way ahead of its time. In 1970, there was relatively little other published knowledge of the plant ecology of Tasmania, and, thus, little basis for determining conservation priorities for species and vegetation types, in a decade of expansion of conservation reserve systems in all parts of Australia, and elsewhere in the world. The literature that existed consisted of broad overviews of the vegetation (Jackson 1965, 1968), a body of work on the dynamics of tall eucalypt forests (Gilbert 1959; Cremer and Mount 1965) and a small number of descriptions, of varying depth, of the vegetation of small parts of the State (e.g. Gibbs 1920; Martin 1940; Davis 1941; Curtis and Sommerville 1947; Gillham 1965). I opportunistically added to this small body of descriptions, selecting some out of the many places that no-one had studied (e.g. Kirkpatrick 1973, 1975; 3.1.4; 3.1.5). Many of the areas I described were almost immediately made into conservation reserves, on the basis of a scientific description being available. It was obvious to me that scientific description was a poor selection criterion for reserves for nature conservation, given the almost total lack of knowledge of the vegetation and flora of most of the State. I thought that the collection of vegetation and species distribution data on a State-wide basis would be necessary to make possible an ecologically appropriate selection of reserves. The provision and analysis of these data became a major focus of my research.

Since 1977, when the first such study was published (Kirkpatrick 1977), State-wide surveys, based on quadrats or releves, have covered all the major vegetation types of Tasmania. The surveys in which I was principal investigator, or supervisor, were of coastal heath (Kirkpatrick 1977; Kirkpatrick and Harris 1999), lentic wetlands (3.5.1; Kirkpatrick and Harwood 1983; Kirkpatrick and Tyler 1988), alpine vegetation (3.1.1; 7.4; Kirkpatrick 1997), grasslands and grassy woodlands (7.5; Kirkpatrick *et al.* 1988; McDougall and Kirkpatrick 1994; Kirkpatrick *et al.* 1995), wet eucalypt forest

(Kirkpatrick *et al.* 1988), *Sphagnum* bogs (Whinam *et al.* 1989), coastal vegetation (Kirkpatrick and Harris 1995), rock plate vegetation (Gilfedder *et al.* 1997) and riparian vegetation (3.5.2). A synopsis of this work, and that conducted by others, to the mid 1990s, was published in Kirkpatrick *et al.* (1995) and its influence is discussed in paper 7.1. I was also involved in the first State-wide survey of coastal sage scrub in California (3.4.1 and 3.4.2).

The only vegetation map of Tasmania was small scale and differentiated only the major formations (Davies 1964). A map better suited to conservation planning was needed. I was involved in the creation of a 1:500,000 vegetation map for the State (Kirkpatrick and Dickinson 1984), and later developed a 1:25,000 vegetation mapping technique (7.6) which has been used to map the whole of the Tasmanian Wilderness World Heritage Area. The field work for the State-wide mapping was somewhat depressing, in that large areas of native vegetation were in the process of clearance, or had just been cleared. I took advantage of the availability of LANDSAT images from 1972 onwards to document the scale and patterning of this clearing (6.1.1 and 6.1.2).

After the pioneering work of Brown *et al.* (1977), I began organizing a data base of the occurrence of vascular plant species in the Tasmanian reserve system, in order to develop priorities for further reservation (Brown *et al.* 1983; Kirkpatrick *et al.* 1991; 7.7). I was later involved in work to document the distribution and reservation status of bryophytes in Tasmania (Moscal and Kirkpatrick 1997; Moscal *et al.* 1997). In the process of undertaking this work it became apparent that there was a need for work on the conservation ecology of threatened plants. This became a major focus of work in my research group from the late 1980s onwards (for a selection of papers, see 4). Many of the threatened species were largely confined to vegetation remnants in the most cleared and developed parts of the State. I therefore decided that it was important to understand how the attributes of remnants affected their biological and conservation attributes. Work to date has been done on vascular plants, bryophytes, lizards and birds (see 5 and 6.1.5 to 6.1.8). Different biotic groups have proven to respond very differently to the characteristics of remnants, and there proved to be no consistent relationship between the

condition of the stands and their importance for threatened elements of biodiversity. For most species, the geometric characters of stands were irrelevant to their survival, and management was critical.

Management was also obviously important for the conservation of non-remnant vegetation. Work on the impacts of trampling by people revealed globally low thresholds for damage in Tasmanian alpine and subantarctic ecosystems (6.5.2 and 6.5.4). The impacts of trampling by horses were shown to vary between different alpine vegetation types (6.5.3). In the first major study of its kind, the impacts of disposal, by burial, of human wastes in the wild country were investigated (6.5.5 and 6.5.6), indicating minimal damage to the vegetation, but waste persistence for at least two years in high rainfall, acid environments.

The alpine vegetation of Tasmania was also very much influenced by fire. Recovery from a single incidence of fire was shown to be exceedingly slow (6.3.1 and 6.3.2), as was recovery from the combined effects of past sheep grazing and fire (6.4.1 to 6.4.3). Fire management was obviously critical in conserving the World Heritage qualities (Kirkpatrick *et al.* 1993) of Tasmanian alpine ecosystems. Most fires started in lowland sedgeland and heaths, making an understanding of their fire management critical in the conservation of alpine vegetation (6.3.4).

Our work on rare or threatened species, and on remnants, and earlier work comparing roadsides and adjacent bush (Fensham and Kirkpatrick 1989), had suggested that vertebrate grazing was playing a major role in determining the species composition and native species richness of lowland grassy vegetation, as well as high country vegetation. The big gaps in understanding were the relationship between grazing regime and vegetation, and the interactions of fire, grazing and vegetation. These were investigated using experimental manipulation and observation (6.4.6 and 6.4.7).

The woodchip export industry, established in the early 1970s in Tasmania, resulted in a massive expansion of logging activity in Tasmania, and a change from selective logging

to clearfelling in the dry eucalypt forests of the State. This was controversial (6.2.1). Research projects on the dynamics of montane eucalypt forests (3.2.4 to 3.2.7) and the comparative impacts on natural vegetation of clearfelling and selective logging (6.2.2 and 6.2.3) were initiated to determine some of the conservation consequences of this silvicultural change.

While most of my research has been in native vegetation outside cities, I was the first researcher in Australia to become interested in the ways in which native ecosystems responded to envelopment by city and suburbs (see 5), and among the first in Australia to take an interest in the plant ecology of the suburban garden (5.7).

Research directed towards improving the conservation of nature in Tasmania required some methodological innovation. I developed the first technique to optimize the selection of reserves (Kirkpatrick *et al.* 1980; 7.3), an achievement recognized by those who have further developed these selection processes (Pressey 2002; Williams *et al.* 2004). I also developed the first technique to quantify wilderness value and calculate its potential loss to development (Kirkpatrick 1979; 7.2), and, with Louise Mendel, devised a method for quantifying natural aesthetic value (7.12). In other methodological research, M.J. Brown and I made the first comparison of the effectiveness of environmental domain techniques and incomplete biological data as inputs to reservation planning (7.9) and M. Fowler and I developed a technique to locate Last Glacial refugia (7.10).

Nature conservation outcomes result from political and social processes as much as scientific understanding. Several of the papers I have included in this volume describe and analyse these processes (1.3, 5.7, 6.1.3, 6.1.4, 6.5.1, 7.1, 7.11).

Pure biogeographical and ecological research can be important for nature conservation either by providing interesting stories that bolster a conservation case, or by increasing understanding of patterns and processes that later turn out to be important in conservation planning or management. Section 2 of the thesis contains floristic biogeographic papers, the earliest of which (2.1) was on alpine floras in Australia. In doing the analysis for this

paper I discovered that the commonly used similarity measures, Sorenson and Jaccard, did not produce reasonable classification and ordination outcomes when some mountain floras were subsets of the flora of larger mountains. I therefore used the number of species in common/the number of species that could be in common. This, and paper 2.4, established that there was a floristic continuum from the oligotrophic superhumid mountains of far southwestern Tasmania to the more fertile, drier and less glaciated mountains of northeast Tasmania and mainland Australia, with most of the variation being in Tasmania. An analysis of the biogeography of vascular plant endemism in Tasmania (2.2) showed the same continuum of variation and also emphasized the role of dolerite, a rock uncommon on mainland Australia. Another paper (2.3) revealed strong centres of local endemism, centres that seemed consistent with a refugium hypothesis, and which were therefore highly important for conservation planning. A graph in paper 2.1 showed a strong relationship between the percentage of endemic species and rainfall, but with the southernmost floras in a line parallel to, and below, the northernmost floras. This led to the investigation reported in paper 2.5, in which strategic rainfall data collection and remote sensing were used to establish that the precipitation maps on which the figure was based were incorrect, with the mountains in the far south receiving the highest precipitation in the State. The technique described in this paper can be adapted to any other areas lacking a reasonable spread of meteorological stations.

Much of my ecological research has been in high mountain and high latitude environments (see the selection of papers in 3.1 and Kirkpatrick 1997). The work in Tasmania contributed substantially to the case for the listing of Tasmanian Wilderness World Heritage Area, demonstrating the globally unusual nature of Tasmanian alpine ecosystems, a distinctiveness relating to a highly maritime climate, oligotrophic soils and isolation (Balmer *et al.* 2004). Several phenomena were researched in detail, for example: the dynamics at different temporal scales of ecosystems dominated by cushion plants; the nature and cause of a sharp floristic boundary between alpine and sedgeland vegetation in southwestern Tasmania; the characteristics and dynamics of unusual fjeldmark vegetation; the ecology of the two species in the genus *Athrotaxis*; the dynamics of

vegetation with *Sphagnum*; changes in the vegetation of the Australian Subantarctic islands.

Other ecological work has been on eucalypt-dominated vegetation (see the selection of papers in 3.2), tropical vegetation in Fiji and the Northern Territory (see the selection of papers in 3.3), Californian vegetation (see the selection of papers in 3.4), wetland vegetation (e.g. 3.5.1 and 3.5.2) and coastal vegetation (e.g. 3.5.4 and 3.5.5). With my students and other colleagues, I have undertaken at least some work on all major terrestrial Tasmanian vegetation types. My work on eucalypts contributed to my major role in the development of the eucalypt-dominated vegetation theme for World Heritage assessment. This theme contributed to the expansion of the listing of the Tasmanian Wilderness in 1989 and the later listing of the Blue Mountains, and played a major role in an influential natural heritage assessment of the Australian Alps (Kirkpatrick 1994).

REFERENCES

Balmer, J., Whinam, J., Kelman, J., Kirkpatrick, J.B. and Lazarus, E., 2004. A review of the floristic values of the Tasmanian Wilderness World Heritage Area. Nature Conservation Report 2004/3, Department of Primary Industries, Water and Environment, Hobart.

Barber, H.N., 1955. Selection in natural populations. *Heredity* 20, 551-572.

Barber, H.N. and Jackson, W.D., 1957. Natural selection in action in *Eucalyptus*. *Nature* 179, 1267-1269.

Brown, M.J., Kirkpatrick, J.B. and Moscal, A., 1983. *An Atlas of Tasmania's Endemic Flora*. Tasmanian Conservation Trust, Hobart.

Cremer, A.W. and Mount, A.B., 1965. Early stages of plant succession following the complete felling and burning of *Eucalyptus regnans* forest in the Florentine Valley, Tasmania. *Australian Journal of Botany* 13, 303-322.

Curtis, W.M. and Sommerville, J., 1947. Boomer Marsh – a preliminary botanical and historical survey. *Papers and Proceedings of the Royal Society of Tasmania* (1947), 151-157.

Davies, G.L., 1964. A vegetation map of Tasmania. *Geographical Review* 54, 249-253.

Davis, C., 1941. Preliminary survey of vegetation near New Harbour, southwest Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* 40, 1-10.

Fensham, R.J. and Kirkpatrick, J.B., 1989. The conservation of original vegetation remnants in the Midlands, Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* 123, 229-46.

Gibbs, L.S., 1920. Notes on the phytogeography and flora of the mountain summit plateaux of Tasmania. *Journal of Ecology* 8, 1-17.

Gilbert, J.M., 1959. Forest succession in the Florentine Valley, Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* 93, 129-151.

Gilfedder, L., Kirkpatrick, J.B. and Ziegeler, D., 1997. Characteristics and conservation status of rock pavement vegetation in lowland perhumid and subhumid Tasmania. A report to the Australian Heritage Commission from the University of Tasmania.

Gillham, M.E., 1965. The Fisher Island field station IV. Vegetation additions and changes. *Papers and Proceedings of the Royal Society of Tasmania* 99, 71-80.

Jackson, W.D., 1965. Vegetation. In J.R. Davies (ed.) *Atlas of Tasmania*. Lands and Survey Department, Tasmania, pp. 30-35.

Jackson, W.D., 1968. Fire, air, water and earth – an elemental ecology of Tasmania. *Proceedings of the Ecological Society of Australia* 3, 9-16.

Kirkpatrick, J.B., 1973. The vegetation of Sloping Island, Tasmania. *Victorian Naturalist* 90, 312-321.

Kirkpatrick, J.B., 1975. Phytosociological analysis of the vegetation of Lagoon Beach, Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* 109, 53-63.

Kirkpatrick, J.B., 1977. *The Disappearing Heath*. Tasmanian Conservation Trust, Hobart.

Kirkpatrick, J.B., 1979. *Hydro-Electric Development and Wilderness in Tasmania*. Department of Environment, Hobart.

Kirkpatrick, J.B., 1994. The international significance of the natural values of the Australian Alps. A report to the Australian Alps Liaison Committee from Tasque.

Kirkpatrick, J.B., 1997. *Alpine Tasmania: An Illustrated Guide to the Flora and Vegetation*. Oxford University Press, Melbourne.

Kirkpatrick, J.B., Barker, P., Brown, M.J., Harris, S. and Mackie, R., 1995. The reservation status of Tasmanian vascular plant communities. Parks and Wildlife Service, Tasmania, Wildlife Scientific Report 95/4.

Kirkpatrick, J.B., Brown, M.J. and Moscal, A., 1980. *Threatened Plants of the Tasmanian Central East Coast*. Tasmanian Conservation Trust, Hobart.

Kirkpatrick, J.B. and Dickinson, K.J.M., 1984. *Vegetation Map of Tasmania*, 1:500,000. Forestry Commission, Tasmania.

Kirkpatrick, J.B., Gilfedder, L. and Fensham, R., 1988. *City Parks and Cemeteries: Tasmania's Grassy Heritage*. Tasmanian Conservation Trust, Hobart.

Kirkpatrick, J.B., Gilfedder, L., Hickie, J. and Harris, S., 1991. Reservation and conservation status of Tasmanian native higher plants. Department of Parks, Wildlife and Heritage, Wildlife Division Scientific Report 91/2.

Kirkpatrick, J.B. and Harris, S. 1995. The conservation of Tasmanian dry coastal vascular plant communities. Parks and Wildlife Service, Tasmania, Wildlife Scientific Report 95/1.

Kirkpatrick, J.B. and Harris, S., 1999. *The Disappearing Heath Revisited*. Tasmanian Environment Centre, Hobart.

Kirkpatrick, J.B. and Harwood, C.E., 1983. The conservation of Tasmanian wetland macrophytic species and communities. *Papers and Proceedings of the Royal Society of Tasmania* 117, 5-20.

Kirkpatrick, J.B., McDougall, K. and Hyde, M., 1995. *Australia's most threatened ecosystem - the southeastern lowland native grasslands*. Surrey-Beatty/WWFA, Sydney.

Kirkpatrick, J.B., Peacock, R.J., Cullen, P.J. and Neyland, M.G., 1988. *The wet eucalypt forests of Tasmania*. Tasmanian Conservation Trust, Hobart.

Kirkpatrick, J.B., Hutchinson, M.N. and MacQuillan, P.B., 1993. Alpine ecosystems. In S. Smith and M.R. Banks (ed). *Tasmanian Wilderness -World Heritage Values*. Royal Society of Tasmania, Hobart, 73-79.

Kirkpatrick, J.B. and Tyler, P. 1988. Tasmanian wetlands and their conservation. In A. MacComb and P.S. Lake (eds.). *The Conservation of Australian Wetlands*. Surrey Beatty and Sons, Sydney, pp. 1-16.

McDougall, K. and Kirkpatrick, J.B. (eds.), 1994. *Conservation of lowland native grasslands in south-eastern Australia*. World Wide Fund for Nature, Australia, Sydney. Martin 1940

Moscal, A. and Kirkpatrick, J.B. 1997. Atlas of mosses and liverworts in Tasmania. A report to the Australian Heritage Commission from the Tasmanian Conservation Trust.

Moscal, A., Askey-Doran, M.J., Kirkpatrick, J.B., Lambourne, M.J., Dalton, P.J. and Seppelt, R., 1997. A preliminary assessment of the conservation and reservation status of Tasmanian bryophytes. A report from the Parks and Wildlife Service for the Regional Forest Agreement Process.

Pressey, R.L., 2002. The first reserve selection algorithm – a retrospective on Jamie Kirkpatrick's 1983 paper. *Progress in Physical Geography* 26, 434-431.

Whinam, J., Eberhard, S., Kirkpatrick, J.B. and Moscal, A., 1989. *Ecology and Conservation of Tasmanian Sphagnum Peatlands*. Tasmanian Conservation Trust, Hobart.

Williams, J.C., ReVelle, C.S. and Levin, S.A., 2004. Using mathematical optimization models to design nature reserves. *Frontiers in Ecology and Environment* 2, 98-105.